



DOI: 10.4274/gulhane.galenos.2024.37267
Gulhane Med J 2025;67(1):8-14

Mapping and spatial analysis of hypertension and diabetes prevalence in selected rural South African communities

© Ushotanefe Useh¹, © Abdullahi Adejare^{1,2}, © Mukadas Akindele^{1,3}, © Sammy Bett⁴

¹North-West University Faculty of Health Sciences, Lifestyle Diseases, Mahikeng, South Africa

²University of Lagos College of Medicine, Faculty of Basic Medical Sciences, Department of Physiology, Lagos, Nigeria; North-West University, Lifestyle Diseases Research Entity, Mahikeng, South Africa

³Bayero University Faculty of Medicine, Department of Physiotherapy, Kano, Nigeria; North-West University, Lifestyle Diseases Research Entity, Mahikeng, South Africa

⁴North-West University, School of Geo and Spatial Science, Department of Geography and Environmental Sciences, Mahikeng, South Africa

Cite this article as: Useh U, Adejare A, Akindele M, Bett S. Mapping and spatial analysis of hypertension and diabetes prevalence in selected rural South African communities. *Gulhane Med J.* 2025;67(1):8-14.

Date submitted:

12.12.2023

Date accepted:

06.08.2024

Epub:

24.12.2024

Publication Date:

20.02.2025

Corresponding Author:

Ushotanefe Useh, PhD, Prof.,
North-West University Faculty of
Health Sciences, Lifestyle Diseases,
Mahikeng, South Africa
+27183892531
22839968@nwu.ac.za

ORCID:

orcid.org/0000-0002-5075-7138

Keywords: Mapping, hypertension, diabetes mellitus, Global Positioning System, medical geography, Kernel density estimation

ABSTRACT

Aims: The widespread prevalence of non-communicable diseases in rural South African communities is concerning. This study aimed to determine the number of cases of hypertension and diabetes reported in rural clinics during the period under review.

Methods: A cross-sectional population-based survey was conducted to determine the prevalence of hypertension and diabetes and the geographical distribution of healthcare facilities in rural communities in the North-West province. Hypertension and diabetes were defined as blood pressure $\geq 130/80$ mm Hg and fasting blood glucose ≥ 126 mg/dL. The range of values for each district municipality (DM) were used to represent map densities and the number of patients screened for hypertension or diabetes.

Results: Hypertension and diabetes map densities were highest in the Bojanala DM and lowest in Dr. Ruth Segomotsi Mompati DM. The number of patients screened for hypertension was highest in the Dr. Ruth Segomotsi Mompati DM (11955-14940) and lowest in the Ngala Modiri Molema DM (11-2996). The number of patients screened for diabetes was highest in the Bojanala DM (12868-16083) and lowest in the Dr. Ruth Segomotsi Mompati DM and Dr. Kenneth Kaunda DM (1-3218).

Conclusions: Although higher numbers of hypertension and diabetes were reported in urban areas, the numbers were grossly low in rural areas. This difference may be associated with the lifestyle of the residents of each area and the uneven distribution of clinics and health centers for the diagnosis and management of these conditions.



Introduction

Worldwide, approximately 70% of all morbidities and mortality are caused by non-communicable diseases (NCDs) (1,2). These NCDs are major risk factors for stroke, myocardial infarction, blindness, and several other cardiovascular diseases (3). A report by the World Health Organization showed that more than 30% of the entire world population suffers from hypertension, whereas more than 10% suffers from diabetes (4). Unfortunately, the large numbers of morbidities and mortality due to NCDs are borne by low- and middle-income countries, where resources are scarce (2). Some reports indicate that approximately 75% of the people in low-and-middle-income countries will have hypertension or other NCDs by the year 2025 (5,6). Among the older population in most African countries, the prevalence is as high as 60%, and a higher prevalence has been reported in urban settings than in rural settings (7). For diabetes, the prevalence is documented as 67-77% in most parts of South Africa (8). Importantly, a study also reported a comorbidity association between hypertension and diabetes (9). This number is increasing (10) and mortality could increase by 2025 (11).

Studies have indicated the possible devastating impacts of the renewed health and behavioral transitions occurring in South Africa (12,13), which has led to an astronomical increase in the prevalence of hypertension, diabetes, and other NCDs (14). Health transitions have been attributed to a demographic shift from rural to urban centers caused by rural-urban migration (15,16). This rural-urban migration leads to dietary and lifestyle modifications and potentiates the population to develop lifestyle diseases (17,18). These transitions have been reported to change the dynamics of the prevalence of hypertension and diabetes and the use of available clinics in the provinces, typifying the situation in the North-West province of South Africa (19).

Medical geography is a relatively new field of study that characterizes the association between pathological and geographical factors that regulate the evolution of pathogens (20). Its use is still limited and uncommon in South Africa and most parts of Sub-Saharan Africa. Medical geography helps map and carry out spatial analyses of diseases and their spread within a locality, aiming to identify low-risk and high-risk regions of a disease (21). For example, in Luxemburg, a study reported geographical variations in the prevalence of hypertension with the highest odds ratio in the industrialized region of the country (22). This study aimed to determine the geographical presentation of hypertension and diabetes concerning the availability of public and private healthcare facilities in rural communities in the North-West Province of South Africa.

Methods

Study design

This was a cross-sectional population-based survey conducted at health clinics in the North-West province of South Africa between April 2015 and March 2016 to determine the distribution of health services in selected rural communities and the number of patients with hypertension and diabetes. There are four districts in the province. At each location, information was gathered using a data collection form detailing the number of clinics, the number of patients diagnosed with hypertension and diabetes, and any pertinent data. Hypertension and diabetes were defined as blood pressure $\geq 130/80$ mmHg and fasting blood glucose ≥ 126 mg/dL respectively (23,24). Additionally, a Global Positioning System receiver was used to collect the geographical coordinates for each center in the study area. These coordinates were then linked to the attribute information collected at the centers and a spatial database containing health center locations and their characteristics was created.

Study locations

The survey was performed in eighty-eight local public and private healthcare facilities in the province. The research team visited 38, 26, 17, and 7 healthcare facilities in Bojanala, Kenneth Kaunda, Ngala Modiri Molema, and Dr. Ruth Segomotsi Mompati district municipalities (DM), respectively.

Dataset

Provincial names and boundaries, as well as their longitudes and latitudes, form the main dataset used in this study. Data were entered into GIS ArcMap and merged with clinical data analyses into one database using common codes for mapping and visualization. Geostatistical spatial analysis was employed to predict the spatial variation in hypertension and diabetes in the province (25).

Inverse distance weighting

Interpolation using inverse distance weighting (IDW) operates under the explicit assumption that objects closer to one another share more similarities than those farther apart (26). With evenly distributed points in an area, the IDW method is effective. In this study, greater weight was assigned to the points closest to the target location; thus, the allocated weights change as an inverse function of " p^{th} power of distance", where the power function (p) is a positive real number. The greater the values closest to the point to be interpolated, the greater the influence. The product of "allotted weights" and "measured values" for all sites was added to provide parameter prediction for the target location (26).

Hotspots analysis

Hotspot Analysis is an analytical technique that combines statistical and spatial methods to measure the concentration of specific elements or attributes. Concerning spatial interpolation, the Hotspot Analysis tool from ArcGIS calculates the Getis-Ord G_i^* statistic for individual features in the dataset (27). The generated z scores and p values were used to identify high and low clustered values in space. This process assesses each feature while considering its context with neighboring features. A feature with a high value may be interesting but not necessarily be a statistically significant hotspot. For a feature to qualify as a statistically significant hotspot, it must have a high value and the surrounding features must possess similarly high values provided all points collected are equally dispersed and at equal intervals (27).

Kriging

Kriging is a stochastic geostatistical technique like inverse distance weighted averaging, where a linear combination of weights at known points is used to predict the value at unknown points. The kriging system is expressed as covariances, commonly derived by estimating and modeling a semi-variogram: a measure of the spatial correlation between two points (28). Kriging is a family of estimators used to interpolate spatial data, including ordinary, universal, and indicator kriging, co-kriging, and others. The choice of kriging depends on the data characteristics and desired spatial model. This study used ordinary kriging to predict unobserved values from observations obtained from nearby clinics. Ordinary kriging is the most common kriging type used in spatial data simulations, and it is considered the best because it minimizes the variance of the estimation error and is more accurate when the unobserved value is closer to the observed value (28).

Statistical Analysis

The range of values for each DM was used to represent map densities and the number of patients screened for hypertension or diabetes.

Ethics approval

This study was conducted following the guidelines for human experiments at North-West University (NWU), South Africa. This study is part of a larger study titled: NCDs, physical activities, and quality of life across different populations in the North-West province. The protocol for this study was approved by the NWU Ethics Committee (NWU-HREC) (registration number: NWU-00014-12-A9, date: 08.03.2012). All procedures for this study were carried out with strict adherence to the National Institutes of Health guidelines for human research.

Results

Density map of the North-West province districts for patients with hypertension and diabetes

The outcome is a less concentrated depiction of a phenomenon that is not portrayed through a sequence of distinct points but through an uninterrupted surface. The density map was found to be highest at the city centers of Bojanala District Municipality (BDM), Dr. Kenneth Kaunda District Municipality (DKKDM), and Ngala Modiri Molema District Municipality (NMMDM). This indicates the frequency of reported cases of hypertension compared with the available number of clinics in the more rural municipalities. There appeared to be a general reduction in density from city centers to suburbs. The density map was generally low at Dr. Ruth Segomotsi Mompati DM (DRSMDM). The Kernel density estimation in Figures 1a and 1b takes the value of the data assigned to a specific point and




Figure 1. Density map for patients with (a) hypertension and (b) diabetes

spreads across a predefined boundary of the northeastern province. This density pattern was also observed in patients with diabetes, with higher concentrations observed in BDM, DKKDM, and NMMDM and lowest concentrations observed in DRSMMDM. This density pattern indicates a possible link between the two studied NCDs.

Number of patients screened for hypertension and diabetes at various clinics in the districts

The study also reported the total number of patients who were screened for hypertension and diabetes at each district. The number of patients screened for hypertension was highest at the Dr. Ruth Segomotsi Mompoti DM (11955-14940) and lowest at the Ngala Modiri Molema DM (11-2996). The number of patients screened for diabetes was highest at the Bojanala DM (12868-16083) and lowest at the Dr. Ruth Segomotsi Mompoti and Dr. Kenneth Kaunda diabetes clinic (1-3218). This is illustrated in Figure 2a, 2b.

Hotspot map showing the prevalence of hypertension and diabetes

Figures 3a and 3b illustrate the hotspot map of the prevalence of hypertension and diabetes in the districts. The pattern was similar for hypertension and diabetes, with the prevalence highest at DKKDM, followed by BDM, DRSMMDM, and finally NMMDM. The map of these hotspots also serves as another pointer to the possible link in the distribution of hypertension and diabetes in the district and the higher presence of these NCDs at urban centers.

Discussion

This study aimed to determine the geographical presentation of hypertension and diabetes in rural communities in the North-West province of South Africa. While a strategic plan to prevent the ever-increasing incidence and prevalence of NCDs like hypertension and diabetes had been developed more than




Figure 2. a, b) Number of patients screened for (a) hypertension and (b) diabetes at various clinics in the districts




Figure 3. a, b) Hotspot map indicating the prevalence of (a) hypertension and (b) diabetes

a decade ago (29) in South Africa, the continuous increase appears to show a lack of understanding of the exact distribution of the affected patients and possible imbalance in the number of healthcare facilities and providers available to manage the conditions. This is the situation in the North-West province, and it explains why there is renewed interest in the likely factors responsible for the mismatch in the province (30). This study presents the observed uneven distribution of patients with either hypertension or diabetes, concerning the number of clinics in the North-West province of South Africa.

In this study, we observed that the density map for hypertension and diabetes showed higher concentrations of the two conditions in the relatively urban centers of Bojanala, Dr. Kenneth Kaunda, and Ngala Modiri Molema DM. It should be noted that these centers have the highest number of clinics in urban areas. The observed high densities could be due to the high population within the major towns in the different districts, as well as pull factors that contribute to the health risks associated with lifestyle diseases. This geographical distribution could also be due to the recently reported lifestyle transition characterized by increased adoption of urban and western lifestyles (7). Other authors have attributed this distribution to dietary transitions (6,7). The traditional diet has been abandoned because of urbanization and rural-urban migration for almost four decades (31). Thus, there appears to be a gradual shift from rural to urban centers (4,5,32), which typifies the situation in the province (7). A close look at the geographical distribution also revealed a high level of similarity between the density map for hypertension and that for diabetes in the entire province.

Importantly, even with fewer clinics in Dr. Ruth Segomotsi Mompoti DM, more people were screened for hypertension than for diabetes. In contrast to this observation, however, the number of patients screened for both hypertension and diabetes at the Bojanala DM mirrors the available number of healthcare facilities. This implies that while the high number of cases reported in the Bojanala district could be due to the available number of clinics, the high number of patients screened for hypertension specifically at the Dr. Ruth Segomotsi Mompoti district may be due to a clear-cut high number of cases in the district. Coincidentally, the number of patients screened for hypertension and diabetes was generally reduced at the Ngala Modiri Molema and Kenneth Kaunda DM compared with the available facilities. One major factor that could affect this distribution is the disparity in the level of education and wealth of the people in the province (33). Well-informed and educated people in major towns tend to have regular medical check-ups at public and private clinics (34). However, their counterparts in rural areas may prefer to use traditional herbs for treatment, as earlier studies have reported extensive use of herbal products by South Africans (35). A higher prevalence of hypertension and diabetes was reported in the Dr. Kenneth Kaunda DM, followed

by Bojanala, Dr. Ruth Segomotsi, and Ngala Modiri Molema DMs. These are the core city centers dominated by educated people whose level of knowledge could directly affect the observed distribution (36,37) and where interprofessional care is commonly practiced in the management of conditions (38).

Based on these findings, there is an urgent need for a more pragmatic approach to managing the astronomical increase in the number of people with NCDs in the province. There may be a need to create policies to regulate nutritional and behavioral lifestyles at city centers while increasing the number of test centers and creating more awareness at rural centers to control the observed geographical distribution of hypertension and diabetes. However, it should be noted that most NCD data were collected and presented using surveying procedures, which might lead to under- or overestimation and are only applicable on coarse spatial scales (39).

This is one of the first studies in the Southern African region that directly used medical geography knowledge to describe a local challenge. The maps provide a real-time graphical representation of hypertension and diabetes in the province.

This study only reported the density maps, and the number of cases observed in the DM of hypertension and diabetes, among several other NCDs like cancer, chronic lung disease, and stroke. The nutritional and behavioral transition in the province is likely to affect the geographical distribution of these other NCDs. In addition, the study only presented the distribution within a short period of just one year. It would have been better if a report covering at least five years was available to provide a clearer picture of the distribution of NCDs in the province. Nonetheless, the report provides insight into the distribution of the two NCDs during the study period.

Conclusion

While higher numbers of hypertension and diabetes were reported in urban areas, the numbers were grossly low in rural areas. This difference may be associated with the lifestyle of the residents of each region and the uneven distribution of clinics and health centers for the diagnosis and management of these conditions.

Acknowledgments

The authors acknowledge the support from the Provincial Department of Health for their efforts in the collection of data for the study.

Ethics

Ethics Committee Approval: The protocol for this study was approved by the NWU Ethics Committee (NWU-HREC) (registration number: NWU-00014-12-A9, date: 08.03.2012).

Informed Consent: Consent form was filled out by all participants.

Footnotes

Authorship Contributions

Concept: A.A., U.U., M.A., Design: U.U., M.A., Data Collection or Processing: U.U., M.A., Analysis or Interpretation: A.A., U.U., S.B., Literature Search: A.A., U.U., Writing: A.A., U.U., S.B., M.A.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2095-2128.
- Mensah GA, Roth GA, Sampson UK, et al. Mortality from cardiovascular diseases in sub-Saharan Africa, 1990-2013: a systematic analysis of data from the Global Burden of Disease Study 2013. *Cardiovasc J Afr*. 2015;26:6-10.
- Stokes J, Kannel WB, Wolf PA, D'Agostino RB, Cupples LA. Blood pressure as a risk factor for cardiovascular disease. The Framingham Study--30 years of follow-up. *Hypertension*. 1989;13:13-18.
- World Health Organisation (WHO). New data highlight increases in hypertension, diabetes incidence. 2012. WHO, 2016. Global report on diabetes. Geneva. Available from: <https://www.who.int/news/item/16-05-2012-new-data-highlight-increases-in-hypertension-diabetes-incidence>
- Seedat YK. Hypertension in developing nations in sub-Saharan Africa. *J Hum Hypertens*. 2000;14:739-747.
- Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Res Clin Pract*. 2011;94:311-321.
- Kaze AD, Schutte AE, Erqou S, Kengne AP, Echouffo-Tcheugui JB. Prevalence of hypertension in older people in Africa: a systematic review and meta-analysis. *J Hypertens*. 2017;35:1345-1352.
- Amod A, Ascott-Evans BH, Berg GI, et al. The 2012 SEMDSA guideline for the management of type 2 diabetes. *J Endocrinol Metab Diabetes S Afr*. 2012;17:61-62.
- Ismail AMA, Saif HFAEA, Taha MM. Effect of alternate nostril breathing exercise on autonomic functions, ocular hypertension, and quality of life in elderly with systemic hypertension and high-tension primary open-angle glaucoma. *Geriatr Nurs*. 2023;52:91-97.
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004;27:1047-1053.
- Bailey SL, Ayles H, Beyers N, et al. Diabetes mellitus in Zambia and the Western Cape province of South Africa: Prevalence, risk factors, diagnosis and management. *Diabetes Res Clin Pract*. 2016;118:1-11.
- Lloyd-Sherlock P, Beard J, Minicuci N, Ebrahim S, Chatterji S. Hypertension among older adults in low- and middle-income countries: prevalence, awareness and control. *Int J Epidemiol*. 2014;43:116-128.
- Akindele MO, Useh U. Chronic diseases of lifestyle risk factor profiles of a South African rural community. *J Public Health Afr*. 2021;12:1006.
- Cheung BM, Li C. Diabetes and hypertension: is there a common metabolic pathway? *Curr Atheroscler Rep*. 2012;14:160-166.
- Gelderblom D, Kok PC. Urbanisation. South Africa's challenge. Vol. 1. Dynamics. Pretoria: HSRC Publishers 1994.
- Solomons NW, Gross R. Urban nutrition in developing countries. *Nutr Rev*. 1995;53:90-95.
- Popkin BM. Nutritional patterns and transitions. *Popul Dev*. 1993;19:138-157.
- MacIntyre UE, Kruger HS, Venterb CS, Vorsterb HH. Dietary intakes of an African population in different stages of transition in the North-West Province, South Africa: the THUSA study. *Nutrition Research*. 2002;22:239-256.
- Kandala NB, Nnanatu CC, Dukhi N, Sewpaul R, Davids A, Reddy SP. Mapping the burden of hypertension in South Africa: a comparative analysis of the National 2012 SANHANES and the 2016 demographic and health survey. *Int J Environ Res Public Health*. 2021;18:5445.
- May JM. Medical geography: its methods and objectives. *Geographical Review*. 1950;40:9-41.
- Rytkönen MJ. Not all maps are equal: GIS and spatial analysis in epidemiology. *Int J Circumpolar Health*. 2004;63:9-24.
- Ruiz-Castell M, Kandala NB, Kuemmerle A, et al. Hypertension burden in Luxembourg: Individual risk factors and geographic variations, 2013 to 2015 European Health Examination Survey. *Medicine (Baltimore)*. 2016;95:e4758.
- Unger T, Borghi C, Charchar F, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75:1334-1357.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2014;37(Suppl 1):81-90.
- Chang KT. Introduction to Geographical Information Systems, McGraw-Hall co.Inc. 2010;2-7.
- Li L, Losser T, Yorke C, Piltner R. Fast inverse distance weighting-based spatiotemporal interpolation: a web-based application of interpolating daily fine particulate matter PM2.5 in the contiguous U.S. using parallel programming and k-d tree. *Int J Environ Res Public Health*. 2014;11:9101-9141.
- Defar A, B Okwaraji Y, Tigabu Z, Persson LÅ, Alemu K. Spatial distribution of common childhood illnesses, healthcare utilisation and associated factors in Ethiopia: evidence from 2016 Ethiopian demographic and health survey. *PLoS One*. 2023;18:e0281606.
- Yamamoto JK. Comparing ordinary kriging interpolation variance and indicator kriging conditional variance for assessing uncertainties at unsampled locations, In: Application of Computers and Operations Research in the Mineral Industry - Dessureault, Ganguli, Kecojevic, & Dwyer editors, Balkema;2005;265-269.
- National Department of Health (NDoH), Statistics South Africa (Stats SA), South African Medical Research Council (SAMRC), ICF. South Africa Demographic and Health Survey 2016:

- Key Indicator Report. Pretoria, South Africa and Rockville, Maryland, USA: NDoH, Stats SA, SAMRC and ICF, 2017.
30. Shisana O, Labadarios D, Rehle T, et al. South African National Health and Nutrition Examination Survey (SANHANES-1). Cape Town:HSRC Press, 2013. Available from: [https://www.hsrc.ac.za/uploads/pageNews/72/SANHANES-launch%20edition%20\(online%20version\).pdf](https://www.hsrc.ac.za/uploads/pageNews/72/SANHANES-launch%20edition%20(online%20version).pdf)
 31. Burkitt DP. Western diseases and their emergence related to diet. *S Afr Med J.* 1982;61:1013-1015.
 32. Kandala N-B, Manda SO, Tigbe WW, Mwambi H, Stranges S. Geographic distribution of cardiovascular comorbidities in South Africa: a national cross-sectional analysis. *J Appl Stats.* 2014;41:1203-1216.
 33. Steyn K. Hypertension in South Africa, Chronic Diseases of Lifestyle in South Africa since 1995-2005. In: Author J, editor. *Chronic Diseases in South Africa.* Chapter 8. 2006. Available from: <https://www.samrc.ac.za/sites/default/files/attachments/2022-10/cdl1995-2005.pdf>
 34. Wreford JT, Esser M, Hippler S. Facilitating relationships between African traditional healing and Western Medicine in South Africa in the time of AIDS: a case study from the Western Cape. CSSR Working Paper 170. Centre for Social Science Research, University of Cape Town: Cape Town. 2006. Available from: <https://open.uct.ac.za/server/api/core/bitstreams/837a8cf7-d38c-4638-9fc0-bf4cfec204bb/content>
 35. Zuma T, Wight D, Rochat T, Moshabela M. The role of traditional health practitioners in Rural KwaZulu-Natal, South Africa: generic or mode specific? *BMC Complement Altern Med.* 2016;16:304.
 36. Hughes GD, Aboyade OM, Clark BL, Puoane TR. The prevalence of traditional herbal medicine use among hypertensives living in South African communities. *BMC Complement Altern Med.* 2013;13:38.
 37. Davids D, Blouws T, Aboyade O, et al. Traditional health practitioners' perceptions, herbal treatment and management of HIV and related opportunistic infections. *J Ethnobiol Ethnomed.* 2014;10:77.
 38. Adejare A, Useh U. Effectiveness of interprofessional care on hypertension in low- and medium-income countries of Africa: a systematic review. *AJBR [Internet].* 2024;27:1-12.
 39. Kauhl B, Maier W, Schweikart J, Keste A, Moskwyn M. Exploring the small-scale spatial distribution of hypertension and its association to area deprivation based on health insurance claims in Northeastern Germany. *BMC Public Health.* 2018;18:121.